Table 1. Comparison of Ultrasound Features in Cervical Cancer Stages IA1–IB2 and Normal Cervical Tissue

| Ultrasound Feature | Normal Cervix | Stage IA1–IB2 Cervical Cancer |
|---|---|---|
| Cervix Shape | Symmetrical, Cylindrical | Asymmetrical, Irregular |
| Cervix Size | Normal Size | Enlarged |
| Cervix Volume | Normal Volume | Increased |
| Cervical Stroma Echogenicity | Homogenous, Hyperechoic | Heterogenous, Hypoechoic |
| Cervical Stroma Vascularity | Normal Vascularity | Increased Vascularity |
| Cervical Stroma Thickness | Normal Thickness | Increased Thickness |
| Cervical Stroma Depth | Normal Depth | Increased Depth |
| Cervical Stroma Invasion | Absent | Present |
| Cervical Canal Obliteration | Absent | Present |
| Cervical Canal Dilatation | Absent | Present |
| Cervical Canal Echogenicity | Hyperechoic | Hypoechoic |
| Cervical Canal Vascularity | Normal Vascularity | Increased Vascularity |
| Cervical Canal Thickness | Normal Thickness | Increased Thickness |
| Cervical Canal Depth | Normal Depth | Increased Depth |
| Cervical Canal